



Multilepton Higgs Decays through the Dark Portal

Tzu Chiang Yuan (阮自強)

Institute of Physics, Academia Sinica




Presented at 广州中山大学物理科学与工程技术学院

第九届TeV 物理工作组学术研讨会

(2014年5月15-18日)


Standard Model/Theory

- SM is a gauge theory
- BEH particle responsible to spontaneously electroweak symmetry breaking



- Provide masses to quarks, leptons, and gauge bosons

Quarks



Leptons

Forces



Outline

- Is that all? Perhaps not.
Exotic Higgs decay might be important.
- Dark U(1) Sector
 - Local Dark U(1)
- Results of 2 and 4 lepton jets
- Summary

Based on Chia-Feng Chang, Ernest Ma, TCY, JHEP 1403 (2014) 054 [arXiv:1308.6071]

Original Higgs U(1) Model As Dark Sector

Chia-Feng Chang, Ernest Ma and TCY, [arXiv:1308.6071]

- The Model

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4}\vec{W}_{\mu\nu} \cdot \vec{W}^{\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu} - \frac{1}{4}C_{\mu\nu}C^{\mu\nu} - \frac{\epsilon}{4}B_{\mu\nu}C^{\mu\nu},$$

Kinetic Mixing

$$\mathcal{L}_{\text{scalar}} = |D_{\mu}\Phi|^2 + |D_{\mu}\chi|^2 - V_{\text{scalar}}(\Phi, \chi),$$

$$V_{\text{scalar}} = -\mu_{\Phi}^2\Phi^{\dagger}\Phi + \lambda_{\Phi}(\Phi^{\dagger}\Phi)^2 - \mu_{\chi}^2\chi^*\chi + \lambda_{\chi}(\chi^*\chi)^2 + \lambda_{\Phi\chi}(\Phi^{\dagger}\Phi)(\chi^*\chi).$$

$$\Phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix}, \quad \chi(x) = \frac{1}{\sqrt{2}}(v_D + h_D(x))$$

- SM Higgs and Dark Higgs Mixing

$$\begin{pmatrix} h_1 \\ h_2 \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} h \\ h_D \end{pmatrix}, \quad \sin 2\alpha = \frac{2m_{12}^2}{m_1^2 - m_2^2}.$$

Heavier state identifies as SM 126 GeV Higgs

Partial List of References

- [1] B. Holdom, PLB166 (1986) 196, PLB178 (1986)65
- [2] Cheung and Yuan, hep-ph/0701107
- [3] Feldman, Liu and Nath, hep-ph/0702123
- [4] Gopalakrishna, Jung and Wells, 0801.3456
- [5] Falkowski, Ruderman, Volansky and Zupan, 1007.3496, 1002.2952
- [6] Davoudiasl, Lee and Marciano, 1203.2947
- [7] Davoudiasl, Lee, Lewis and Marciano, 1304.4935
-

Rich Higgs Phenomenology

- Higgs mixings implies non-standard Higgs decay

$$h_1 \rightarrow \gamma_D \gamma_D$$


$$h_1 \rightarrow h_2 h_2$$

$$h_1 \rightarrow h_2 h_2^* \rightarrow h_2 \gamma_D \gamma_D$$

$$h_1 \rightarrow h_2 h_2 h_2$$

- h_2 decays predominantly into two dark photons $h_2 \rightarrow \gamma_D \gamma_D$
- Dark photon may decay to light SM leptons through mixings (kinetic mixing with photon or Stueckelberg mass terms) $\gamma_D \rightarrow \bar{l} l \quad (l = e, \mu)$
- This model predicts: non-standard modes of Higgs decay could be 4, 8, or even 12 leptons in the final states!
- Similar signals (with MET) can occur in hidden valley scenarios, SUSY, Weinberg's Higgs Portal model etc.

Kinematic Region ($m_1 = 126$ GeV)



Higgs Decay Width

- SM Higgs total width contains two pieces

$$\Gamma_{h_1} = \cos^2 \alpha \hat{\Gamma}_h + \Gamma_{h_1}^{NS},$$

- Non-standard contributions from dark Higgs U(1) sector


$$\Gamma_{h_1}^{NS} = \sin^2 \alpha \hat{\Gamma}(h_1 \rightarrow \gamma_D \gamma_D) + \Gamma(h_1 \rightarrow h_2 h_2) + \Gamma(h_1 \rightarrow h_2 \gamma_D \gamma_D) + \Gamma(h_1 \rightarrow h_2 h_2 h_2) + \dots$$

- The SM Higgs width is 4.03 MeV (Theory) while LHC constrains the non-standard Higgs width to be less than 3.3 MeV (See talk by 陳國明)

Parameter Space: $\alpha, g_D, m_2, m_{\gamma_D}$



Contour Plots for non-standard branching ratio for standard model Higgs ($\sin^2\alpha=0.0009; Br=0.1$)

Chia-Feng Chang, Ernest Ma and TCY, [arXiv:1308.6071]



Light Mass Region

Contour Plots for non-standard branching ratio for standard model Higgs ($\sin^2\alpha=0.0009; Br=0.2$)



Light Mass Region

Multilepton Jets at the LHC

- Four processes which may lead to signals of multilepton jets at the LHC



- (I) and (II) are SM processes
- (III) and (IV) are dark processes
- (III) with $X=Z$ is SM process with modified h_1 coupling

- (I) $pp \rightarrow h \rightarrow ZZ \rightarrow l^+l^-l^+l^-$
 - (II) $pp \rightarrow VV \rightarrow l^+l^-l^+l^-$ ($VV = ZZ, \gamma\gamma, Z\gamma$)
 - (III) $pp \rightarrow h_1 \rightarrow XX \rightarrow l^+l^-l^+l^-$ ($XX = ZZ, \gamma_D\gamma_D, h_2h_2$)
 - (IV) $pp \rightarrow h_1 \rightarrow h_2h_2 \rightarrow \gamma_D\gamma_D\gamma_D\gamma_D \rightarrow l^+l^-l^+l^-l^+l^-l^+l^-$




Figure 6. Some topologies of 4 (left) and 2 (right) lepton-jets for process III. The 4 lepton-jets can also be coming from the SM of process I with h_1 replaced by the SM h . The immediate state of h_2h_2 for the 2 lepton-jets is not shown since the branching ratio for $h_2 \rightarrow l^+l^-$ is very tiny.





Figure 7. Some topologies of 4 (left) and 2 (right) lepton-jets for process IV.

Benchmark Points

Benchmark Point	g_D	M_{γ_D}	m_2	$\text{Br}_{h_1 \rightarrow \text{DarkStuff}}$	$\text{Br}_{h_2 \rightarrow \gamma_D \gamma_D}$	$\text{Br}_{\gamma_D \rightarrow l+l^-}$
A	0.005	1.5	4	$\sim 16\%$	99%	50%
B	0.009	1.8	10	$\sim 20\%$	100%	50%
C	0.005	1.5	40	$\sim 15\%$	99%	50%
D	0.005	1.8	40	$\sim 11\%$	99%	50%

Table 1. Several benchmark points of the dark portal used to calculate the signals of multilepton jets ($\epsilon = 10^{-4}$ and $\sin^2 \alpha = 10^{-3}$). Mass in GeV.

Number of Events vs. M_{Inv}




- Benchmark point B
- No cuts
- Blue strip (I)
- Black dash (II)
- Red solid (III)
- Yellow strip (IV)

Kinematical Cuts

Gopalakrishna, Jung and Wells, 0801.3456;
 Falkowski, Ruderman, Volansky and Zupan, 1007.3496;
 ATLAS collab. 1302.4403.

Basic cuts: (4 leptons case) $p_{T_i} \geq 20, 10, 10, 10 \text{ GeV}$, $|\eta| < 2.3$;
 (8 leptons case) $p_{T_i} \geq 20, 10, 10, 10, 0, 0, 0, 0 \text{ GeV}$, $|\eta| < 2.3$,

4 lepton-jets cuts : $\Delta R_{j_i j_j}^d > 0.7$, $\Delta R_{l_i l_j}^s < 0.2$, $M_{\text{Invariant}} = M_{h_1} \pm 10 \text{ GeV}$;
2 lepton-jets cuts : $\Delta R_{j_1 j_2}^d > 0.7$, $\Delta R_{l_i l_j}^s < 0.2$, $M_{\text{Invariant}} = M_{h_1} \pm 10 \text{ GeV}$.



Cross Sections (fb) at Benchmark Points

- 2 and 4 lepton-jets cuts have great impact on SM processes, in particular for process (I) where $gg \rightarrow h_1 \rightarrow ZZ \rightarrow 4$ leptons compared with process (II) where $qq \rightarrow ZZ \rightarrow 4$ leptons
- Process III where $h_1 \rightarrow \gamma_D \gamma_D \rightarrow 4$ leptons favors by 2 lepton-cuts
- Lepton-jets cuts have non-trivial effects on process IV where $h_1 \rightarrow h_2 h_2 \rightarrow \gamma_D \gamma_D \gamma_D \gamma_D \rightarrow 8$ leptons; in general it has smaller cross section than process III

- (I) $pp \rightarrow h \rightarrow ZZ \rightarrow l^+ l^- l^+ l^-$
- (II) $pp \rightarrow VV \rightarrow l^+ l^- l^+ l^-$ ($VV = ZZ, \gamma\gamma, Z\gamma$)
- (III) $pp \rightarrow h_1 \rightarrow XX \rightarrow l^+ l^- l^+ l^-$ ($XX = ZZ, \gamma_D \gamma_D, h_2 h_2$)
- (IV) $pp \rightarrow h_1 \rightarrow h_2 h_2 \rightarrow \gamma_D \gamma_D \gamma_D \gamma_D \rightarrow l^+ l^- l^+ l^- l^+ l^- l^+ l^-$

Cuts	Benchmark Point	I	II	III	IV
Basic	A	0.118	70.7	95.3	23.2
	B	0.118	70.7	204	45.8
	C	0.118	70.7	96.7	19.2
	D	0.118	70.7	68.3	13.1
Basic + 4 Lepton-Jets	A	9.63×10^{-3}	0.337	9.86×10^{-3}	$\leq 10^{-10}$
	B	9.63×10^{-3}	0.337	9.80×10^{-3}	$\leq 10^{-10}$
	C	9.63×10^{-3}	0.337	9.93×10^{-3}	3.05
	D	9.63×10^{-3}	0.337	9.84×10^{-3}	0.92
Basic + 2 Lepton-Jets	A	$\leq 10^{-10}$	0.08	95.3	1.75
	B	$\leq 10^{-10}$	0.08	201	$\leq 10^{-10}$
	C	$\leq 10^{-10}$	0.08	95.8	$\leq 10^{-10}$
	D	$\leq 10^{-10}$	0.08	68.2	$\leq 10^{-10}$

Table 2. Cross sections (in unit of fb) at the LHC-14 for the background processes (I and II) and dark sector processes (III and IV) with the basic, 4 and 2 lepton-jets cuts at the 4 benchmark points.

Summary

- Exotic Higgs decay modes are important since it may lead to discovery of a hidden sector like a dark $U(1)$
- Multilepton modes should be studied in more details, in particular multi-muon jets (or even multi-pion jets)
- Challenging but very interesting signals with small SM background!